



Uploaded to the VFC Website

▶▶▶▶ 2016 ◀◀◀◀

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](#)

If Veterans don't help Veterans, who will?

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



Antipsychotic drugs more frequently used in Alzheimer's disease patients, new study finds

Published on October 14, 2015 at 8:37 AM

Antipsychotic drugs are initiated in patients with Alzheimer's disease (AD) more frequently than in the general population - already 2-3 years before the Alzheimer's diagnosis, according to a new study from the University of Eastern Finland. Most commonly, antipsychotics were initiated during the six months following the Alzheimer's diagnosis; however, the incidence of new antipsychotic users was high also later on. The results were published in British Journal of Psychiatry.

The study analysed the incidence of antipsychotic use eight years before and four years after the Alzheimer's diagnosis. During the follow-up, one third of the persons with Alzheimer's disease initiated antipsychotics. The incidence of antipsychotic use in persons with Alzheimer's disease was five times more common than in the control group not diagnosed with AD. The study is based on an analysis of 7,217 Finnish patients with Alzheimer's disease, whose use of antipsychotics was investigated as part of the Finnish MEDALZ-2005 Study comprising nation-wide register data.

An earlier study analysing the duration of antipsychotic treatment in the same study population found that the use of antipsychotics in patients with Alzheimer's disease continues for a significantly long period. More than half (57%) were long-term users, i.e. their antipsychotic use lasted more than one year.

According to the Finnish Current Care Guidelines relating to the treatment of memory disorders, the use of antipsychotics should be limited to a short-term treatment of severe psychotic symptoms, agitation and aggression. Furthermore, as the risks of antipsychotics may outweigh their benefits among persons with Alzheimer's disease, many guidelines, for example those issued by the UK-based National Institute for Health and Care Excellence and the American Psychiatric Association recommend that antipsychotics should be used only in treatment of severe psychotic symptoms, aggression and agitation when the symptoms cause significant distress or risk of harm to the patient or others.

"In Finland, the present incidence of antipsychotic use and the duration of antipsychotic treatment are not in line with the treatment guidelines. This is a cause of concern, because the use of antipsychotics increases the risk of severe adverse incidents such as stroke, falling and fracture, and death," says Professor Sirpa Hartikainen of the University of Eastern Finland.

Source:

University of Eastern Finland
