



Uploaded to VFC Website

~ November 2012 ~

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

[Veterans-For-Change](#)

*Veterans-For-Change is a 501(c)(3) Non-Profit Corporation
Tax ID #27-3820181*

If Veteran's don't help Veteran's, who will?

We appreciate all donations to continue to provide information and services to Veterans and their families.

https://www.paypal.com/cgi-bin/webscr?cmd=_s-xclick&hosted_button_id=WGT2M5UTB9A78

Note:

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members.

Item ID Number 03833 ☐ **Not Scanned**

Author Arnold, E. L.

Corporate Author Department of Life and Behavioral Sciences, USAF Aca

Report/Article Title Typescript: A Rapid Method for the Determination of
Several Phenoxyalkanoic Acid Herbicides in Soil
Samples

Journal/Book Title

Year 1974

Month/Day February

Color ☐

Number of Images 16

Description Notes Includes two memorandums clearing the typescript for
publication

DEPARTMENT OF THE AIR FORCE
HEADQUARTERS UNITED STATES AIR FORCE ACADEMY
USAF ACADEMY, COLORADO 80840



REPLY TO
ATTN OF: OIPM (Capt Boyle/4050)
SUBJECT: Review of Manuscript

20 FEB 1974

TO: DFLS (Capt Young)

The Department of Defense has reviewed your manuscript, "A Rapid Method for the Determination of Several Phenoxyalkanoic Acid Herbicides in Soil Samples," co-authored by Maj. E.L. Arnold. It is cleared, with no changes, for open publication.

William F. Frensley
WILLIAM F. FRENSLEY, Major, USAF
Deputy Director of Information

Atch
Manuscript (2)

Cy to: DFSS



"MAN'S FLIGHT THROUGH LIFE IS SUSTAINED BY THE POWER OF HIS KNOWLEDGE"

DEPARTMENT OF THE AIR FORCE
HEADQUARTERS UNITED STATES AIR FORCE
WASHINGTON, D.C.



REPLY TO
ATTN OF:

SAF/OIS (74-103)

15 February 1974

SUBJECT:

Paper, "A Rapid Method for the Determination of Several
Phenoxyalkanoic Acid Herbicides in Soil Samples"

TO:

AFA/OI

The Department of Defense has reviewed subject paper and
has no objection to open presentation.

FOR THE CHIEF OF STAFF

A handwritten signature in cursive script, reading "John J. Pelszynski", is written over the typed name.

JOHN J. PELSZYNSKI
Colonel, USAF
Chief, Office for Security Review
Office of Information

1 Atch
Subj as above

A RAPID METHOD FOR THE DETERMINATION OF SEVERAL PHENOXYALKANOIC
ACID HERBICIDES IN SOIL SAMPLES

CLEARED
FOR OPEN PUBLICATION

FEB 13 1974 20

DIRECTORATE FOR SECURITY REVIEW (OASD-PA)
DEPARTMENT OF DEFENSE

by

E. L. ARNOLD AND A.L. YOUNG

DEPARTMENT OF LIFE AND BEHAVIORAL SCIENCES

USAF ACADEMY, COLORADO 80840

90400
740103

BRIEF

A rapid gas chromatographic method for the measurement of several phenoxyalkanoic acid herbicides in soil samples is described. Average recovery of herbicides added to soil ranged from 99.7% to 87.5% ($\pm 4.9\%$).

INTRODUCTION:

The problems associated with the ecologically safe disposal of contaminated or excess pesticides ^{have} ~~has~~ received increased attention in recent years. Among those compounds requiring a safe method for disposal are the halogenated phenoxyalkanoic acid herbicides, 2,4-dichlorophenoxyacetic acid (2,4-D), 2,4,5 trichlorophenoxyacetic acid ^(2,4,5-T) ~~(2,3,5-T)~~, and their esters. One of the means of disposal of these herbicides which has been suggested is soil biodegradation, whereby the herbicides are incorporated in soil and degraded by soil microorganisms. (1). In order to evaluate the effectiveness of this technique, analytical procedures are necessary for the measurement over a wide range of concentrations of both the n-butyl esters and the sodium salts of 2,4,-D and 2,4,5-T. Since evaluation of the biodegradation technique will require the analysis of large numbers of soil samples taken from relatively large field plots, an analytical procedure which is not only accurate and precise but also rapid and simple to perform is required. A number of different techniques have been devised for this type of analysis (2-4), however, many of these procedures tend to be complex and time consuming since they were designed for samples where herbicide concentrations were low and many interfering substances were present. This paper will describe a simple, rapid and accurate procedure for the simultaneous determination of the n-butyl esters of 2,4-D and 2,4,5-T and their corresponding hydrolysis products in soil samples.

EXPERIMENTAL:

APPARATUS: A Beckman Gas Chromatograph, Model GC-45 equipped with a hydrogen flame ionization detector was employed. Columns were 6 ft x 2 mm i.d. glass u-tubes packed with 80/100 mesh chromasorb W coated with 10% DC-200 silicone gum as the stationary phase. All columns used were supplied by the instrument manufacturer. Conditions for the chromatography were: temperatures; inlet, 245°C; column, 220°C; detector, 250°C; air flow: carrier (helium), 60 cc/min; detector make up (helium), 60 cc/min; hydrogen, 50cc/min; air, 250 cc/min. One microliter samples were injected directly on column.

REAGENTS: Methanol, chloroform, acetylchloride and anthracene were ACS reagent grade chemicals (J.T. Baker) and were used without additional purification. The sodium salts of 2,4 dichlorophenoxyacetic acid (2,4-D) and 2,4,5 trichlorophenoxyacetic acid (2,4,5-T) were practical grade (Dow Chemical Co.). The n-butyl esters of 2,4-D and 2,4,5-T were analyzed to be greater than 99% purity (Hercules, Inc.): 3% solution of anhydrous hydrogen chloride in methanol was prepared by reacting 5 ml of acetyl chloride with 100 ml of methanol. (5).

PROCEDURES: Standard solutions of the n-butyl esters and the sodium salts of 2,4-D and 2,4,5-T were prepared by dissolving weighed amounts of each compound in either chloroform (n-butyl esters) or methanol (sodium salts). After serial dilutions were made, the sodium salt solutions were reacted with a 3% (w/w) solution of anhydrous HCl in methanol (5 ml herbicide salt standard solution plus 1.5 ml 3% HCl: methanol) in order to form methyl esters suitable for chromatography. After heating for 15 minutes at 55°C, a sufficient volume of a chloroform solution of anthracene internal standard was added to obtain a

final internal standard concentration of 10 $\mu\text{g/ml}$, 50 $\mu\text{g/ml}$ or 250 $\mu\text{g/ml}$ depending upon the herbicide concentration of the standard solution. Sufficient internal standard solution was also added to the n-butyl ester solutions to obtain the same range of internal standard concentrations as used with the sodium salts. All standard solution volumes were adjusted to 10 ml with chloroform prior to chromatography.

Standard curves for each herbicide were prepared by the chromatography of standard solutions of varying herbicide concentration, calculating the ratios of the individual herbicide GC peak areas to the GC peak area of the internal standard and plotting these ratios against the actual herbicide concentrations in the standard solutions. Concentration curves were prepared for each of the four herbicides at three different concentration ranges, (0-10 $\mu\text{g/ml}$, 10-100 $\mu\text{g/ml}$, 100-1000 $\mu\text{g/ml}$).

Soil samples used for evaluating the procedures were prepared by dissolving weighed amounts of each of the four herbicides in anhydrous methanol and adding this solution to representative control soil samples contained in 1 liter round bottom flasks. Solvent was then removed from the samples using a rotary flash evaporator. Control soil samples had been previously analyzed and shown to be free from herbicide residues.

Soil samples were analyzed in the following manner. Five grams of treated soil were placed in 15 ml glass screw top culture tubes, and the herbicides extracted with 10 ml anhydrous methanol by heating the sealed tubes at 55°C in a water bath. Samples were shaken periodically during the heating period. After cooling, a 5 ml aliquot of the methanol extract was transferred to a clean culture tube and mixed with 1.5 ml of 3% HCl:methanol.

Reaction was allowed to proceed at 55°C for 15 minutes. The samples were allowed to cool to room temperature and ~~3.5~~^{3.5} ml of a chloroform solution of internal standard added. The final concentration of internal standard obtained was either 10[?] g/ml, 50[?] g/ml or 250[?] g/ml, depending upon the expected range of herbicide present. One microliter samples were analyzed by gas chromatography. Soil concentrations were determined by multiplying the concentrations in mg/ml obtained from the standard curves by a factor of four to obtain the concentrations in micrograms of each herbicide per gram of soil. Soil samples obtained from biodegradation test plots were analyzed in the same manner. pH say
mg/ml

RESULTS AND DISCUSSION:

Employing the chromatographic conditions outlined above, the four herbicides of interest could be separated from each other in less than ten minutes as shown in Figure I. This figure also illustrates the freedom of the samples from interfering substances which might be present in the soil. In the chromatogram, the herbicides which were present in the soil as either salts or free acids appear as methyl esters. The esterification reaction appeared to produce a quantitative yield of methyl esters regardless of the equilibrium form of the herbicide which was present.

Five replicate soil samples for each of five different herbicide concentrations were analyzed. All samples were analyzed on the same day as they were prepared in order to minimize hydrolysis of n-butyl esters due to the prevailing alkalinity (pH 8.4) of the soil which was used. The results of these analysis are given in Table I. Three additional replicates at two concentration levels were stored at 4°C for 24 hrs then analyzed for the purpose of evaluating the degree of hydrolysis. These data are presented

TABLE I

RECOVERIES OF HERBICIDES ADDED TO SOIL SAMPLES*

Compound	Amount added mg/g soil	Average Recovered** (mg/g)	Percentage Recovery
2,4-D, Na	2.00	2.19	110
2,4,-D, n-butyl	2.00	1.85	93
2,4,5-T, Na	2.00	2.08	104
2,4,5-T, n-butyl	2.00	1.84	92
2,4-D, Na	1.00	1.12	112
2,4-D, n-butyl	1.00	0.86	86
2,4,5-T, Na	1.00	0.98	98
2,4,5-T, n-butyl	1.00	0.88	88
2,4-D, Na	0.250	0.264	106
2,4-D, n-butyl	0.250	0.212	85
2,4,5-T, Na	0.250	0.251	100
2,4,5-T, n-butyl	0.250	0.201	81
2,4-D, Na	0.050	0.061	112
2,4-D, n-butyl	0.050	0.048	96
2,4,5-T, Na	0.050	0.052	104
2,4,5-T, n-butyl	0.050	0.041	82
2,4-D, Na	0.010	0.012	120
2,4-D, n-butyl	0.010	0.007	70
2,4,5-T, Na	0.010	0.009	90
2,3,4-T, n-butyl	0.010	0.007	70

* Samples analyzed immediately following preparation.

**Mean value of 6 analyses.

in Table II. Average recovery of total added herbicide ranged from 99.7% at soil concentration of 8 mg/g to 87.5% at soil concentrations of 0.04 mg/g. The mean relative standard deviation for all four compounds of interest was $\pm 4.9\%$. It appears from the data contained in Table I that approximately 8% of the n-butyl esters of 2,4,5-T and 12% of the n-butyl esters of 2,4-D contained in any sample are converted to methyl esters by the derivative forming reaction. Since this rate of conversion is relatively constant at all herbicide concentrations, a conversion factor may be added in order to improve the accuracy of the analysis. The data presented in Table II indicates a relatively rapid hydrolysis of the n-butyl esters of 2,4-D and 2,4,5-T when exposed to alkaline soil. In the 24 hr period of exposure, 9% of the n-butyl ester of 2,4-D and 8% of the n-butyl ester of 2,4,5-T were hydrolyzed. It should be noted that in this calculation it is presumed that some of the hydrolysis of esters noted, is due to the derivative forming reaction. This presumption is based upon studies made with different concentrations of HCl. At catalyst concentrations lower than 3% (w/w), recovery of the n-butyl esters from soil samples approached 100% while recovery of the herbicide salts decreased rapidly if the samples were analyzed immediately after preparation. These data illustrate the necessity of analyzing field soil samples as soon after removal from the test plots as possible if a comparison of esters and salts is necessary.

At the conclusion of the analysis of the prepared soil samples, a number of samples were obtained from USAF soil biodegradation test plots and analyzed for herbicide residues. A herbicide formulation of the n-butyl esters of 2,4-D and 2,4,5-T had been incorporated at several different application rates into the soil of these plots. Table III contains the data

TABLE II

HYDROLYSIS OF HERBICIDE ESTERS ADD TO ALKALINE SOIL SAMPLES

Compound	Amount added mg/g soil	Time of exposure hrs	Average amount recovered	Percent increase/decrease of 24 hr exposure
2,4-D, Na	2.00	0	2.19	
2,4-D, Na	2.00	24	2.71	+26.0
2,4-D, n-butyl	2.00	0	1.85	
2,4-D, n-butyl	2.00	24	1.38	-24.0
2,4,5-T, Na	2.00	0	2.08	
2,4,5-T, Na	2.00	24	2.40	+16.0
2,4,5-T, n-butyl	2.00	0	1.84	
2,4,5-T, n-butyl	2.00	24	1.58	-13.0
2,4-D, Na	1.00	0	1.12	
2,4-D, Na	1.00	24	1.32	+20.0
2,4-D, n-butyl	1.00	0	.86	
2,4-D, n-butyl	1.00	24	.72	-14.0
2,4,5-T, Na	1.00	0	0.98	
2,4,5-T, Na	1.00	24	1.14	+16.0
2,4,5-T, n-butyl	1.00	0	0.88	
2,4,5-T, n-butyl	1.00	24	0.71	-19.0

TABLE III

DETERMINATION OF HERBICIDE CONCENTRATIONS IN SOIL BIODEGRADATION TEST PLOTS

Sample #	Soil Depth	2,4-D Na mg/g	2,4-D n-butyl mg/g	2,4,5-T Na mg/g	2,4,5-T n-butyl mg/g
1-1	0-6"	0.740	0.376	0.700	0.326
1-2	"	0.698	0.354	0.650	0.303
1-3	"	0.731	0.368	0.684	0.340
1-4	"	0.740	0.385	0.706	0.327
1-5	"	0.726	0.361	0.671	0.316
Mean $\pm \sigma$		0.727 \pm .018	0.369 \pm .012	0.682 \pm .023	0.322 \pm .014
2-1	0-6"	0.835	1.05	0.410	1.62
2-2	"	0.910	1.15	0.426	1.70
2-3	"	0.856	1.08	0.410	1.66
2-4	"	0.795	1.06	0.392	1.57
2-5	"	0.852	1.10	0.408	1.60
Mean $\pm \sigma$		0.850 \pm .042	1.09 \pm .040	0.409 \pm .010	1.63 \pm .043
3-1	6-12"	0.125	0.015	0.098	0.032
3-2	"	0.130	0.015	0.100	0.042
3-3	"	0.117	0.013	0.110	0.035
3-4	"	0.115	0.011	0.095	0.040
3-5	"	0.132	0.014	0.112	0.038
Mean $\pm \sigma$		0.124 \pm .015	0.013 \pm .002	0.103 \pm .008	0.037 \pm .004

obtained from five replicate analyses from three different samples taken from the test plots. As can be seen from the data in the table, the concentrations of the different herbicide residues in these plots varied greatly, however, the mean relative standard deviation ($\pm 6.1\%$) for all samples varied only slightly from that determined when prepared samples were used. This indicates that the analytical technique is applicable for the determination of herbicide residues over the large concentration ranges found in actual field samples.

In addition to the relatively high precision and accuracy of the analytical procedure, one of its major advantages is the number of samples which can be analyzed for four different components in a short period of time. The time required for one individual to analyze 12 soil samples was 3-4 hours or about 30 minutes per sample. Sensitivity of the procedure allows accurate measurement of any of the four compounds at levels greater than 4 microgram/gram soil. It was also noted during the investigation that, if desired, two other ester formulations (octyl and iso-octyl) of 2,4-D and 2,4,5-T could be determined simultaneously with the analysis described previously.

ACKNOWLEDGEMENT:

The authors gratefully acknowledge Anthony M. Wachinski for providing the soil samples from soil biodegradation test plots used in this work. Further reproduction is authorized to satisfy the needs of the U.S. Government.

LITERATURE CITED

1. M.A. Loos, Degradation of Herbicides, P.C. Kearney and D.D. Kaufman, Eds., Marcel Dekker, Inc., New York, 1969, pp 18-24.
2. D.E. Clark, J. Agr. Food Chem., 17:1168 (1969).
3. D.E. Gutnick and G. Zweig, J. Chromatog. 13:319 (1964).
4. M.L. Taylor, R.L.C. Wu, E.L. Arnold, B.M. Hughes and T.O. Tiernan, Abstracts; 21st Annual Conference of the American Society for Mass Spectrometry, San Francisco, Calif., May 1973.
5. Applied Science Laboratories, Inc., State College, Pa., Catalog 16, p 78, 1973.

List of Figure Captions

Figure 1: Gas chromatogram of the extract of a prepared soil sample containing 1 mg/g each 2,4-D and 2,3,5-T sodium salts and 1 mg/g each 2,4-D and 2,4,5-T n-butyl esters. Salts were converted to methyl esters prior to chromatography. Chromatographic conditions as outlined in experimental section, attenuation = 1000x.

RECORDER RESPONSE

