

Uploaded to the VFC Website

▶ ▶ 2016 ◀ ◀

This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

Veterans-For-Change

If Veterans don't help Veterans, who will?

Note: VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



Research: Adenosine deaminase may help activate immune system against HIV

Published on February 3, 2016 at 4:43 AM

New research findings published in the February 2016 issue of the *Journal of Leukocyte Biology*, suggest that a new therapeutic strategy for HIV may already be available by repurposing an existing prescription drug. The drug, an enzyme called adenosine deaminase, or ADA, ultimately may be able to activate the immune system against HIV and to help the immune system "remember" the virus to prevent or quickly eliminate future infection.

"We hope this study puts ADA in the spotlight as a powerful immune modulator in vaccine strategies enhancing anti-HIV immune responses and limiting the need for life-lasting treatments," said Núria Climent, Ph.D., a researcher involved in the work from the Retrovirology and Viral Immunopathology Laboratory, at AIDS Research Group from IDIBAPS in Barcelona, Spain.

To make their discovery, scientists studied the effects of ADA in cells from HIV patients and non-infected individuals. Dendritic cells derived from blood cells were exposed to HIV-1 proteins or whole inactivated HIV-1 virus in the presence or absence of ADA. The degree of lymphocyte proliferation, regulatory T-cell generation and cytokine secretion were measured. ADA addition resulted in adenosine degradation leading to a reduction of regulatory T-cell mediated suppression. Notably, the presence of ADA produced an increase in CD4+ responder T cells, in CD8+ T cell proliferation and in T cell memory generation. An increase in the secretion of immunologically relevant Th1 cytokines was also seen.

"We need to find new strategies that will empower the immune system towards long-term control of HIV infection," said Luis J. Montaner, D.V.M., M.Sc., D.Phil., Editor-in-Chief of the *Journal of Leukocyte Biology*. "The availability of an approved drug that already targets the mechanisms described here ensures the quick translation of this work from the bench to the clinical."

Source:

Federation of American Societies for Experimental Biology