

Uploaded to the VFC Website



This Document has been provided to you courtesy of Veterans-For-Change!

Feel free to pass to any veteran who might be able to use this information!

For thousands more files like this and hundreds of links to useful information, and hundreds of "Frequently Asked Questions, please go to:

Veterans-For-Change

If Veterans don't help Veterans, who will?

Note

VFC is not liable for source information in this document, it is merely provided as a courtesy to our members & subscribers.



Anti-platelet drug may slow spread of colon and pancreatic cancer cells

Dec 15 2016

Researchers from Oregon Health and Science University have found that aspirin may slow the spread of some types of colon and pancreatic cancer cells. The paper is published in the *American Journal of Physiology—Cell Physiology*.

Platelets are blood cells involved with clotting. They promote the growth of cancerous cells by releasing growth factors and increasing the response of certain proteins that regulate tumor cell development (oncoproteins). Low doses of aspirin, an anti-platelet drug, have been shown to reduce the risk of some types of gastrointestinal cancers, but the process by which aspirin hampers tumor growth has been unclear. "The current study was designed to determine the effect of inhibition of platelet activation and function by aspirin therapy on colon and pancreatic cancer cell proliferation," the researchers wrote.

The research team combined activated platelets primed for the clotting process with three groups of cancer cells:

- metastatic colon cancer (cells that have spread outside the colon),
- nonmetastatic colon cancer (cells that grow only within the colon) and
- nonmetastatic pancreatic cancer cells.

When they added aspirin to the mixture, they found that the platelets were no longer able to stimulate growth and replication in the pancreatic and nonmetastatic colon cancer cells. The metastatic colon cancer cells continued to multiply when treated with aspirin.

In pancreatic cancer cells, low doses of aspirin stopped the platelets from releasing growth factor and hampered the signaling of the oncoproteins that cause cancer to survive and spread. Only very high doses—larger than are possible to take orally—were effective in stopping growth in the metastatic colon cells, explained the researchers.

The findings detail the interaction among platelets, aspirin and tumor cells and are promising for the future treatment of nonmetastatic cancer, according

to the researchers. "Our study reveals important differences and specificities in the mechanism of action of high- and low-dose aspirin in metastatic and nonmetastatic cancer cells with different tumor origins and suggests that the ability of aspirin to prevent platelet-induced c-MYC [an oncoprotein] expression might be selective for a nonmetastatic phenotype."

Source:

American Physiological Society (APS)